arXiv:1108.1305v1 [quant-ph] 5 Aug 2011

Violation of time reversal symmetry in quantum noninvasive measurements

Adam Bednorz,[1, *] Kurt Franke,[2] and Wolfgang Belzig[2]
[1]*Faculty of Physics, University of Warsaw, Hoża 69, PL-00681 Warsaw, Poland*
[2]*Fachbereich Physik, Universität Konstanz, D-78457 Konstanz, Germany*

(Dated: August 8, 2011)

A quantum system in equilibrium is by definition in a state, which is invariant under time-reversal. However, in general projective quantum measurements of incompatible observables would obviously break the time-reversal symmetry, which one could try to circumvent by a weak quantum measurement leaving the state of the system intact. We show that, paradoxically, time-reversal symmetry is violated for this type of measurement, with the violation detectable in higher-order correlation functions. We propose an experiment with quantum dots to measure this apparent violation of time reversal symmetry in a third-order current correlation function.

Microscopically, classical physics is symmetric under time reversal[1, 2]—the arrow of time appears solely due to macroscopic loss of information which is the essence of the second law of thermodynamics. In quantum mechanics time reversal symmetry holds for standard measurements of compatible observables[3]. Strong, projective measurements of incompatible quantities affect future measurements and break the general time symmetry just as invasive classical measurements do, although a residual symmetry (unrelated to the one discussed here) under exchange of pre- and postselected states survives[4]. However, the situation gets complicated when considering noninvasive quantum measurements[5, 6]. In this Letter we demonstrate that time reversal symmetry is broken in general for measurements of incompatible observables even in the weak, non-disturbing limit. We also propose an experimentally feasible test of our conjecture in a quantum transport setup. Assuming experimental confirmation, our result shows that quantum systems with time reversal symmetry will not *appear* to have time reversal symmetry. This is inconsistent with classical intuition and shows that the weak measurement limit in quantum mechanics cannot properly be considered noninvasive.

Time reversal is often presented visually as reversing the direction of a movie[7]. If the reversed film is compatible with the equations of motion, the system is symmetric under time reversal. The intuitiveness of this explanation is complicated by the necessity of making measurements to create the movie. Time reversal symmetry appears to be broken due to the second law of thermodynamics, which provides our arrow of time in the direction of increasing entropy. This bias can be eliminated by considering stationary systems in thermal equilibrium where, for example, a backward-running clock would be just as likely as a forward-running clock.

We consider an abstract "movie" to be the result of a series of measurements. Once the parameters of the system, measured quantities and times of measurements are all specified, the measurement results—the movies—are described by a joint probability density $P(a_1(t_1),\dots,a_n(t_n))$. A probabilistic description is required by the statistical nature of the ensemble, measurement noise, and, in the case of quantum measurements, quantum randomness. The time reversed experiment involves reversed initial conditions (e.g. position $\mathbf{r}^T=\mathbf{r}$ and momentum $\mathbf{p}^T=-\mathbf{p}$), external forces (e.g. magnetic field $\mathbf{B}^T=-\mathbf{B}$), measured quantities $a\to a^T$ (^{T} denotes the time reversal), and time—and therefore, ordering—of the measurements. Assuming for the moment that the measurements have no effect on the time evolution, there will be an equivalence between the probability densities of the forward (P) and reversed (P^T) sets of measurements:

$$P(a_1(t_1),\dots,a_n(t_n))=P^T(a_n^T(-t_n),\dots,a_1^T(-t_1))\ . \tag{1}$$

This result leads to the principle of detailed balance[1] and reciprocity of thermodynamic fluxes [2]. Equation (1) can be tested by comparing moments of the distribution. Of special interest is the moment where each measurement appears linearly:

$$\langle a_1(t_1)\cdots a_n(t_n)\rangle=\langle a_n^T(-t_n)\cdots a_1^T(-t_1)\rangle^T\ . \tag{2}$$

The time reversal operation is here applied to every measured quantity, the parameters of the initial state and the evolution equations. In such a form, (2) is independent of the validity of charge conjugation and parity symmetries and also of relativistic invariance [3, 8, 9]. For low-energy physics, time reversal alone is already a good symmetry in the equations of motion so, for equilibrium systems with no external magnetic field, $P=P^T$.

For equations (1) and (2) to hold, the measurements should not disturb the system, since any disturbance would create an asymmetry between before and after the measurement. On the other hand, quantum measurements of commuting observables clearly satisfy (1), so non-disturbance of the system is too strong a requirement: what matters is whether or not the measurements disturb each other. We adopt a straightforward definition that a set of measurements is not disturbed by some additional operation or perturbation if the probability density for the measurements is not affected. If a measurement can be added or removed without affecting the joint probability density of the remaining measurements, then the complete set of measurements are mutually non-disturbing. In other words, integrating over any single measurement should yield the same distribution that would be obtained if that measurement were never performed,

$$\int \mathrm{d}a_j P(a_1,\dots,a_n) = P(a_1,\dots,\cancel{a}_j,\dots,a_n) \qquad (3)$$

A set of measurements performed on an equilibrium ensemble and satisfying (3) *should therefore satisfy* (1) *and* (2). This principle works not only for non-disturbing classical measurements, but also for quantum measurements of compatible observables (e.g. space-like separated[3]). These latter types of measurements disturb the system but are mutually non-disturbing in the sense of (3). Incompatible projective measurements [10] do not satisfy (3) but also do not in general satisfy (1). Nevertheless, for projective measurements one can find another symmetry—unrelated to (2)—under the exchange of boundary conditions in pre- and postselected ensembles[4, 6].

Weak measurements, introduced by Aharonov, Albert and Vaidman [5], however, allow for mutually non-disturbing measurements of incompatible quantities [11]. They are also useful in refuting macrorealism by violation of the Leggett-Garg inequality [12], which has been recently tested experimentally [13]. In this scheme, the probability describing the measurement outcome P is a convolution $P = D * Q$ of a large white noise D (independent of the measured system, with uncertainty inversely proportional to the strength of the measurement) with a quantum quasi-probability Q – a sometimes negative Wigner-type distribution which is independent of the measurement strength [14]. The construction is based on positive operator-valued measures [15], using Gaussian Kraus operators [16, 17] and taking the limit of infinite uncertainty. For the details, see [18] and the Supplemental Material A. Importantly, the pseudo-probability density for a series of weak measurements satisfies (3). Additionally, weak measurements result in only infinitesimal changes to the density matrix (and therefore, to the entropy) of any single system in the ensemble. Correlations of weak measurements are given by [18, 19]

$$\left\langle \prod_k a_k(t_k) \right\rangle = \mathrm{Tr}\{A_n(t_n), ..\{A_2(t_2), \{A_1(t_1), \rho\}\}..\}/2^n \qquad (4)$$

for $t_n \geq \cdots \geq t_2 \geq t_1$ and $\{A, B\} = AB + BA$. Here $A(t)$ denotes a quantum operator corresponding to the quantity a at time t and ρ is the initial density operator. *The striking paradox is that weak measurements satisfy* (3)*, at least as a limiting procedure, but it can easily be verified that weak measurement correlations* (4) *in general violate the time reversal property* (2). This asymmetry is only present for fast measurements of three or more incompatible observables. In the case of compatible ($AB = BA$) or only two (not necessarily compatible) measurements the ordering is irrelevant and the symmetry (2) holds. For slow measurements, each operator $A(t)$ in (4) is replaced with $\int f(t)A(t)dt$, where $f(t)$ turns on and off slowly compared to relevant timescales of the system. This slow measuring smoothes the resulting distribution Q so that any antisymmetric contributions vanish and therefore (1) and (2) will still apply.



FIG. 1: The double well, described effectively by two states $|l\rangle$ and $|r\rangle$, with energy shift 2ε and tunneling amplitude τ. Even in the ground state, quantum fluctuations allow jumps between the wells, which turn to be not time symmetric under noninvasive measurement.

Let us demonstrate the paradox in a simple system consisting of a particle in a double-well potential as in Fig. 1. The particle is effectively described by the ground states of the left and right wells, $|l\rangle$ and $|r\rangle$ respectively. Higher excited states have much more energy and for low temperatures can be ignored, leaving an effective two state system. Using the basis states, the operator for expected location is $Z = |l\rangle\langle l| - |r\rangle\langle r|$, and the effective Hamiltonian reads

$$H = \varepsilon(|l\rangle\langle l| - |r\rangle\langle r|) + \tau(|l\rangle\langle r| + |r\rangle\langle l|), \qquad (5)$$

where 2ε is the energy difference between wells and τ is the tunneling amplitude. Since no magnetic fields are present, H and Z must be even under time reversal. We are now in a position to test equation (2) with z measured at three separate times and with the initial thermal state $\rho \propto \exp(-H/kT)$. The correlation for three weak measurements can be calculated using (4) and $Z(t) = e^{iHt/\hbar} Z e^{-iHt/\hbar}$:

$$\langle z(t_1)z(t_2)z(t_3)\rangle = \alpha(\varepsilon^2 + \tau^2 \cos(2(t_3 - t_2)\Delta/\hbar)), \qquad (6)$$

where $\Delta = \sqrt{\varepsilon^2 + \tau^2}$, $\alpha = -(\varepsilon/\Delta^3)\tanh(\Delta/kT)$. For this system and measurements, the expression corresponding to the right hand side of (2) differs from (6) only by the exchange of $t_3 - t_2$ with $t_2 - t_1$. However, (6) is clearly asymmetric under this exchange, demonstrating that time reversal symmetry is broken for correlations of quantum weak measurements.

Despite the simplicity of the above example, a genuine, fast weak detection scheme is probably difficult to implement experimentally in this case. Below, we present a



FIG. 2: (color online) (a) The system consisted of the dot (red) exchanging the elementary charge with a reservoir (blue). (b) The diagram of energy levels. At zero temperature, the Fermi sea of blue levels is full but electrons may still jump on and off the dot's red level.(c) Proposed detection by an electric junction (yellow). The junction and dot are weakly coupled capacitively and the directly measured quantity is the current I through the junction biased with the voltage. The charge is allowed to jump between the dot and the lower reservoir but not the junction.

more realistic example, leveraging recent developments in quantum dots [20]. We consider a quantum dot containing a single energy level ε, coupled to a Fermi reservoir by an energy independent coupling described effectively by the tunneling rate $\Gamma/\hbar$, as depicted in Fig. 2a,b. The occupation n on the dot (classically either 0 or 1 in elementary charge units) is the measured observable N. The quantum observable and the Hamiltonian read [21]

$$\begin{aligned} &N = c^\dagger c,\ H = \varepsilon N + \\ &\int dE\, [\sqrt{\Gamma/2\pi}\, c^\dagger\psi(E) + \text{h.c.} + E\psi^\dagger(E)\psi(E)], \end{aligned} \tag{7}$$

which describes energy-independent tunneling between the dot and reservoir, where ε is dot level energy. We assume usual fermion anticommutation relations $\{\psi,\phi\} = 0$, $\{\psi^\dagger,\phi\} = 0$ if $\psi \neq \phi$, $\{c^\dagger, c\} = 1$ and $\{\psi^\dagger(E),\psi(E')\} = \delta(E-E')$. Spin is neglected here but if necessary all results can be simply multiplied by 2. The initial state is $\rho \propto \exp(-H/k_BT)$. The Hamiltonian (7) and the occupation are certainly symmetric under time reversal, $H^T = H$ and $N^T = N$.

To show the time asymmetry we will use the frequency domain, defining the third cumulant

$$S_3^N(\omega,\omega') = \int dt dt'\, e^{i\omega t + i\omega' t'} \langle \delta n(t)\delta n(t')\delta n(0)\rangle \tag{8}$$

with $\delta n = n - \langle n\rangle$. The asymmetry-probing quantity is the imaginary part of the third cumulant $\mathrm{Im}S_3^N(\omega,\omega')$, which should vanish if (2) holds. To calculate (8) we use the close-time-path formalism [22, 23], defining matrices in 2×2 Keldysh space

$$\check N = \begin{pmatrix} 1 & 0 \\ 0 & 1/4 \end{pmatrix},\ \check G = \begin{pmatrix} G^K & G^R \\ G^A & 0 \end{pmatrix} \tag{9}$$

with $G^R(\omega) = i\hbar/(\hbar\omega - \varepsilon + i\Gamma/2) = -G^{A*}(\omega)$ and $G^K(\omega) = \tanh(\hbar\omega/2k_BT)\hbar\Gamma/(2(\hbar\omega-\varepsilon)^2 + \Gamma^2/2)$. Then

$$\begin{aligned} S_3^N(\omega,\omega') = -\int \frac{d\alpha}{2\pi} \mathrm{Tr}\check G(\alpha)\check N \times \\ [\check G(\alpha+\omega) + \check G(\alpha+\omega')]\check N \check G(\alpha+\omega+\omega')\check N. \end{aligned} \tag{10}$$

The integral can be performed analytically but the result contains digamma functions at finite temperatures. As suspected, $\mathrm{Im}S_3^N$ is not zero, see Fig. 3. Both imaginary and real parts vanish far from resonance. The asymmetry is the strongest at low temperatures ($kT \ll \varepsilon$) and for comparable energy, tunneling and frequency scales ($\varepsilon \sim \Gamma \sim \hbar\omega$). This suggests that it is zero-point fluctuations of the charge jumping on and off the dot that are responsible for the asymmetry. The symmetry is restored if one of ω, ω', or $\omega+\omega'$ is equal to 0. As expected, $\mathrm{Im}S_3^N$ vanishes for slow measurements $\omega,\omega' \ll \Gamma/\hbar$. In the limit $\omega,\omega' \to 0$, the result for S_3^N is a special case of the application of full counting statistics [23].

For the experimental confirmation of the asymmetry one must introduce a weakly coupled detector. We propose an electric voltage-biased junction coupled weakly to the dot, so that its conductance depends on the charge on the dot, see Fig. 2c. The externally measured quantity is the current I through the junction, in particular $I \simeq I^0 + \chi N$, where I^0 is the intrinsic current in the junction and χ is its susceptibility due to the dot's charge. Then $S_3^I \simeq S_3^{I0} + \chi^3 S_3^N$ where S_3^{I0} is the internal current noise of the junction in the absence of the dot. Measurements of S_3^{I0} have been demonstrated [24] therefore we expect the measurement of S_3^I to be feasible. Such an experiment will confirm the time reversal symmetry violation only if the dot is not driven out of equilibrium. It is always possible for a certain parameter range, see Supplemental Material B, for the detailed model.

We have shown that the property (3) is not sufficient to guarantee time-reversal symmetries in sequences of quantum measurements. One could argue that the weak measurement still affects the system and forces time direction in this way. On the other hand, one expects a natural limit where the influence on the system is negligible and the time symmetry should hold. Although the entropy of the ensemble of many runs/copies of the systems is indeed changed by a finite value, even in the weak limit, we can safely assume the independence and identity between systems so that the entropy change per system is small.

In conclusion, we have shown that, contrary to classical intuition, quantum noninvasive measurements violate time reversal symmetry. This violation is effectively a



FIG. 3: (color online) Time asymmetry of the third order correlation of dot occupation fluctuations $S_3^N(\omega,\omega')$ in frequency domain at $kT \ll \Gamma$. The symmetry is broken if Im $S_3^N \neq 0$. (a) For $\varepsilon/\Gamma = 0.5$ and arbitrary ω, ω' the asymmetry vanishes (white lines) at when ω, ω' or $\omega+\omega'$ is equal to 0. The hexagonal structure reflects the symmetry under permutations of frequencies. (b) Taking different values $\varepsilon/\Gamma = 1, 0.5, 0.2$ and $\omega = \omega'$ it is clear that the maximal asymmetry occurs for energy/frequency parameters of the same order.

failure of weak measurements to accurately reflect time-reversal symmetry inherent in a system. As such, it is independent of the validity of other symmetries such as charge-parity-time. Since quantum measurements of finite strength manifestly break time reversal invariance, our result shows that, in contrast to classical measurements, all quantum measurements break time reversal invariance regardless of their strength. Finally, we emphasize that only successful realization of the proposed or analogous experiment will ultimately confirm our conjecture.

We are grateful to Bertand Reulet, Christoph Bruder and Tomasz Dietl for helpful discussions. Financial support from the DFG through SFB 767 and SP 1285 is acknowledged.

---

* Electronic address: Adam.Bednorz@fuw.edu.pl

[1] N.G. van Kampen, *Stochastic Processes in Physics and Chemistry*, (North-Holland, Amsterdam, 2007).
[2] L. Onsager, Phys. Rev. **37**, 405 (1931).
[3] R.F. Streater, A.S. Wightman, *PCT, spin and statistics, and all that* (Benjamin, New York, 1964).
[4] Y. Aharonov, P.G. Bergmann, and J.L. Lebowitz, Phys. Rev. **134**, B1410 (1964); M. Gell-Mann, and J. Hartle, in *Physical Origins of Time Asymmetry* (eds. J. Halliwell, J. Perez-Mercader, and W. Zurek) 311 (Cambridge University Press, Cambridge, 1994) arXiv:gr-qc/9304023.
[5] Y. Aharonov, D.Z. Albert, and L. Vaidman, Phys. Rev. Lett. **60**, 1351 (1988).
[6] Y. Aharonov, S. Popescu, and J. Tollaksen, Phys. Today **63**, i.11, 27 (2010).
[7] D. Griffiths, *Introduction to Elementary Particles* (WILEY-VCH, Weinheim, 2008).
[8] O.W. Greenberg, Phys. Rev. Lett. **89**, 231602 (2002).
[9] M. Sozzi, *Discrete Symmetries and CP Violation* (Oxford U.P., New York, 2008)
[10] J. von Neumann, *Mathematical Foundations of Quantum Mechanics* (Princeton U.P., Princeton, 1932).
[11] J.S. Lundeen *et al.*, Nature **474**, 188 (2011)
[12] A.J. Leggett, A. Garg, Phys. Rev. Lett. **54**, 857 (1985).
[13] A. Palacios-Laloy *et al.*, Nature Phys. **6**, 442, (2010).
[14] W.P. Schleich, *Quantum Optics in Phase Space*, (Wiley-VCH, Weinheim, 2001); A.I. Lvovsky *et al.*, Phys. Rev. Lett. **87**, 050402 (2001); D. Bozyigit *et al.*, Nature Phys. **7**, 154 (2011); M. P. da Silva *et al.*, Phys. Rev. A **82**, 043804 (2010); C. Eichler *et al.*, Phys. Rev. Lett. **106**, 220503 (2011).
[15] H. M. Wiseman and G. J. Milburn *Quantum Measurement and Control* (Cambridge University Press, Cambridge, 2009).
[16] K. Kraus, *States, Effects and Operations* (Springer, Berlin, 1983).
[17] A. Barchielli, L. Lanz, and G.M. Prosperi, Nuovo Cimento B **72**, 79 (1982); C.M. Caves, and G.J. Milburn, Phys. Rev. A **36**, 5543 (1987); A. Schmid, Ann. Phys. **173**, 103 (1987).
[18] A. Bednorz, and W. Belzig, Phys. Rev. Lett. **105**, 106803 (2010).
[19] B. Berg *et al.*, Phys. Rev. A **80**, 033624 (2009); M. Tsang, **80**, 033840 (2009); H.F. Hofmann, **81**, 012103 (2010); J. Dressel, S. Agarwal, and A.N. Jordan, Phys. Rev. Lett. **104**, 240401 (2010); K. Chou *et al.*, Phys. Rep. **118**, 1 (1985).
[20] W. Lu *et al.* Nature **423**, 422 (2003); Sukhorukov, E.V. *et al.*, Nature Phys. **3**, 243 (2007).
[21] Y.M. Blanter, and M. Büttiker, Phys. Rep. **336**, 1 (2000).
[22] J. Schwinger, J. Math. Phys. **2**, 407 (1961); L.V. Keldysh, JETP **20**, 1018 (1965); L.P. Kadanoff, G. Baym, *Quantum Statistical Mechanics* (Benjamin, New York, 1962); A. Kamenev, and A. Levchenko, Advances in Phys. **58**, 197 (2009).
[23] Y. Utsumi, Phys. Rev. B **75**, 035333 (2007).
[24] B. Reulet, J. Senzier, and D.E. Prober, Phys. Rev. Lett. **91**, 196601 (2003); Y. Bomze *et al.*, **95**, 176601 (2005); G. Gershon *et al.*, **101**, 016803 (2008); J. Gabelli, and B. Reulet, J. Stat. Mech. P01049 (2009) doi:10.1088/1742-5468/2009/01/P01049.

Supplemental Material



FIG. 4: The idea of weak measurement. The measured system (yellow) instantly interacts with a prepared ancilla (red wavefunction), which is measured projectively afterwards (green detector). The procedure is repeated with identical but independent ancillas.

A. WEAK MEASUREMENT

To justify Eq. (4) in the manuscript we consider a series of weak measurements. Following Aharonov, Albert and Vaidman [5], each weak measurement introduces an ancilla system and creates entanglement via an instant interaction Hamiltonian $\hat{H}_I = \hbar\delta(t) g\hat{p}\hat{A}$ where g is the strength of interaction, $\hat{p}$ is momentum operator of the ancilla, conjugate to position $\hat{x}$ ($[\hat{x},\hat{p}] = i\hbar$), and $\hat{A}$ is the measured observable. The interaction is followed by von Neumann projection [10] of the ancilla onto a position eigenstate which destroys the ancilla. The system can however be measured again with the next ancilla, as shown in Fig. 4. The density matrix after the j^{th} measurement is

$$\hat{\rho}_j = e^{-ig_j\hat{p}_j\hat{A}_j}\left(\hat{\rho}_{j-1}\otimes|\phi_j\rangle\langle\phi_j|\right)e^{ig_j\hat{p}_j\hat{A}_j} \tag{A.1}$$

where $|\phi_j\rangle$ is the initial prepared state of ancilla j. By inserting identity operations $\int da\,|a\rangle\langle a| = \hat{1}$, the measurement interaction can be expressed as shifts of the ancilla wavefunction.

$$\hat{\rho}_j = \int da'_j\, da''_j |\phi_j(x_j - g_j a'_j)\rangle\langle\phi_j(x_j - g_j a''_j)| \times|a'_j\rangle\langle a'_j|\hat{\rho}_{j-1}|a''_j\rangle\langle a''_j| \tag{A.2}$$

In (A.2), the the state of ancilla j which has the shifted wavefunction $\phi_j(x_j - g_j a'_j)$ is written as $|\phi_j(x_j - g_j a'_j)\rangle$. The joint probability $P(a_1,\ldots,a_n) =: P(\boldsymbol{a})$ is the probability of measuring the ancillas in a set of position eigenstates with positions given by $x_k = a_k/g_k$. This scaling makes the resulting distribution independent of measurement strength in both the strong and weak limits.

$$P(\boldsymbol{a}) = \mathrm{Tr}\left\{\hat{\rho}_n\prod_k\frac{1}{g_k}|a_k/g_k\rangle\langle a_k/g_k|\right\} \tag{A.3}$$

$$= \int d\boldsymbol{a'}\, d\boldsymbol{a''}\,\tilde{\rho}_n\left(\boldsymbol{a'},\boldsymbol{a''}\right) \times\prod_k\frac{1}{g_k}\phi\left(g_k(a_k - a'_k)\right)\phi^*\left(g_k(a_k - a''_k)\right) \tag{A.4}$$

In (A.4), $\tilde{\rho}_j$ is defined recursively by

$$\tilde{\rho}_j(a'_1,a''_1,\ldots,a'_j,a''_j) = \langle a'_j|a'_{j-1}\rangle\tilde{\rho}_{j-1}\langle a''_{j-1}|a''_j\rangle. \tag{A.5}$$

Using Gaussian wavefunctions $\phi(x) = (2\pi)^{-1/4}\,e^{-x^2/4}$, a change of variables to $\bar{\boldsymbol{a}} = (\boldsymbol{a'}+\boldsymbol{a''})/2$ and $\boldsymbol{\delta a} = \boldsymbol{a'}-\boldsymbol{a''}$ separates the joint probability density into a quasiprobability signal (Q) and detector noise (D).

$$\begin{aligned} P(\boldsymbol{a}) &= \int d\bar{\boldsymbol{a}}\; D(\boldsymbol{a}-\bar{\boldsymbol{a}})\,Q(\bar{\boldsymbol{a}}),\\ D(\boldsymbol{a}-\bar{\boldsymbol{a}}) &= \prod_k g_k|\phi(g_k(a_k-\bar{a}_k)|^2,\\ Q(\bar{\boldsymbol{a}}) &= \int d\boldsymbol{\delta a}\; e^{-(\boldsymbol{g}\cdot\boldsymbol{\delta a})^2/2}\;\tilde{\rho}_n(\bar{\boldsymbol{a}},\boldsymbol{\delta a}) \end{aligned} \tag{A.6}$$

The measure $d\boldsymbol{\delta a}$ depends on $\bar{\boldsymbol{a}}$. Eq. (A.6) is the joint quasiprobability density for the series of von Neumann measurements. The quasiprobability has a well defined limit $g \to 0$. In this limit for time-resolved measurement, the averages with respect to this quasiprobability are given by

$$\langle a_1\cdots a_n\rangle = \int d\boldsymbol{a}'d\boldsymbol{a}''\,\tilde{\rho}(\boldsymbol{a}',\boldsymbol{a}'')\prod_k\frac{a'_k+a''_k}{2}, \tag{A.7}$$

which is equivalent to (4). The genuine, measured probability $P = Q * D$ is positive definite because it contains also the large detection noise $\sim 1/g$ which is Gaussian, white and completely independent of the system.

An alternative, equivalent approach is based on Gaussian POVMs and special Kraus operators [15-17]. Let us begin with the basic properties of POVM. The Kraus operators $\hat{K}(a)$ for an observable described by $\hat{A}$ with continuous outcome a need only satisfy $\int da\hat{K}^\dagger(a)\hat{K}(a) = \hat{1}$. The act of measurement on the state defined by the density matrix $\hat{\rho}$ results in the new state $\hat{\rho}(a) = \hat{K}(a)\hat{\rho}\hat{K}^\dagger(a)$. The new state yields a normalized and positive definite probability density $P(a) = \mathrm{Tr}\,\hat{\rho}(a)$. The procedure can be repeated recursively for an arbitrary sequence of (not necessarily commuting) operators $\hat{A}_1,\ldots,\hat{A}_n$,

$$\hat{\rho}(a_1,\ldots,a_n) = \hat{K}(a_n)\hat{\rho}(a_1,\ldots,a_{n-1})\hat{K}^\dagger(a_n)\,. \tag{A.8}$$

The corresponding probability density is given by $P(a_1,\dots,a_n) = \mathrm{Tr}\ \hat\rho(a_1,\dots,a_n)$. We now define a family of Kraus operators, namely $\hat K_\lambda(a) = (2\lambda/\pi)^{1/4}\exp(-\lambda(\hat A - a)^2)$. It is clear that $\lambda\to\infty$ should correspond to exact, strong, projective measurement, while $\lambda\to 0$ is a weak measurement and gives a large error. In fact, these Kraus operators are exactly those associated with the von Neumann measurements previously described (with $\lambda = g^2$). We also see that strong projection changes the state (by collapse), while $\lambda\to 0$ gives $\hat\rho(a)\sim\hat\rho$, and hence this case corresponds to weak measurement. However, the repetition of the same measurement k times effectively means one measurement with $\lambda\to k\lambda$ so, with $k\to\infty$, even a weak coupling $\lambda\ll 1$ results in a strong measurement. For an arbitrary sequence of measurements, we can write the final density matrix as the convolution

$$\hat\rho_{\boldsymbol\lambda}(\boldsymbol a) = \int da'\,\hat\varrho_{\boldsymbol\lambda}(\boldsymbol a')\prod_k g_k(a_k - a'_k) \tag{A.9}$$

with $g_k(a) = e^{-2\lambda_k a^2}\sqrt{2\lambda_k/\pi}$. Here $\boldsymbol\lambda = (\lambda_1,\dots,\lambda_n)$ $\boldsymbol a = (a_1,\dots,a_n)$, and $da = da_1\dots da_n$. The quasidensity matrix $\hat\varrho$ is given recursively by

$$\begin{aligned}\hat\varrho_{\boldsymbol\lambda}(\boldsymbol a) \;=\; \int\frac{d\xi}{2\pi}e^{-i\xi a_n}\int\frac{d\phi}{\sqrt{2\pi\lambda_n}}e^{-\phi^2/2\lambda_n}\times\\ e^{i(\xi/2+\phi)\hat A_n}\,\hat\varrho_{\boldsymbol\lambda}(a_1,\dots,a_{n-1})e^{i(\xi/2-\phi)\hat A_n}\end{aligned} \tag{A.10}$$

with the initial density matrix $\hat\varrho = \hat\rho$ for $n = 0$. We can interpret g in (A.9) as some internal noise of the detectors which, in the limit $\lambda\to 0$, should not influence the system. We *define* the quasiprobability [18] $Q_{\boldsymbol\lambda} = \mathrm{Tr}\,\hat\varrho_{\boldsymbol\lambda}$ and abbreviate $Q\equiv Q_{\mathbf 0}$. In this limit (A.10) reduces to

$$\hat\varrho(\boldsymbol a) = \int\frac{d\xi}{2\pi}e^{-i\xi a_n}e^{i\xi\hat A_n/2}\hat\varrho(a_1,\dots,a_{n-1})e^{i\xi\hat A_n/2}\,. \tag{A.11}$$

Note that $Q_{0\dots0,\lambda} = Q$, so the last measurement does not need to be weak (it can be even a projection). The averages with respect to Q are easily calculated by means of the generating function (A.11), e.g. $\langle a\rangle_Q = \mathrm{Tr}\,\hat A\hat\rho$, $\langle ab\rangle_Q = \mathrm{Tr}\,\{\hat A,\hat B\}\hat\rho/2$, $\langle abc\rangle_Q = \mathrm{Tr}\,\hat C\{\hat B,\{\hat A,\hat\rho\}\}/4$ for $\boldsymbol a = (a,b,c)$. As a straightforward generalization to continuous measurement, we obtain

$$\begin{aligned}&\langle a_1(t_1)\cdots a_n(t_n)\rangle_Q = \\ &\mathrm{Tr}\hat\rho\{\hat A_1(t_1),\{\cdots\{\hat A_{n-1}(t_{n-1}),\hat A_n(t_n)\}\cdots\}\}/2^{n-1}\end{aligned} \tag{A.12}$$

for time ordered observables, $t_1\le t_2\le\cdots\le t_n$. Both above presented ways agree in the limit $\lambda = g^2\to 0$.

B. DETECTOR

An effective model of weakly detecting the dot's charge using an electric junction is shown in Fig. 5. The junction is treated as another dot between two reservoirs but in a broad level regime. The complete Hamiltonian, consisting of the dot part (7), and the junction part, reads



FIG. 5: The model of detecting the dot's charge. An electric junction contains another dot with effective occupation N', coupled capacitively to the measured dot. The fluctuations of the current I in the junction biased by the voltage V depend on the dot's occupation N with the proportionality constant χ.

$$\begin{aligned}&\hat H + \varepsilon'\hat N' + \hat H_V + e^2\hat N\hat N'/C + eV\hat Q_L + \int dE\ \times\\ &\sum_{A=L,R}[\sqrt{\Gamma'/2\pi}\hat d^\dagger\hat\psi_A(E) + \mathrm{h.c.} + E\psi_A(E)\hat\psi_A(E)],\\ &\hat N_L = \hat\psi^\dagger_L(E)\psi_L(E),\ \hat N' = \hat d^\dagger\hat d,\end{aligned} \tag{B.1}$$

where $\hat N_L$ is the total number of elementary charges e in the left reservoir, C is the capacitance between the dot and the QPC, Γ',ε' denote effective tunneling rate and level energy of the QPC and V is the bias voltage.

We measure current fluctuations in the junction, $I(t)$, with the current in Heisenberg picture defined as $\hat I(t) = -ed\hat N_L(t)/dt$. Such fluctuations have already been measured experimentally at low and high frequencies [24]. Most of fluctuations are just generated by the shot noise in the junction. Now, we consider a finite, but still very large capacitance. We expect a contribution from the system dot's charge fluctuation to S^I_3 of the order C^{-3}. We assume separation of the system's and detector's characteristic frequency scales, namely

$$(\Gamma,\varepsilon,k_BT)\ll eV\ll(\Gamma',\varepsilon'), \tag{B.2}$$

which also includes the broad level approximation for the detector's dot. There exists a special parameter range,

$$\frac{e^2}{\Gamma' C}\gg\frac{\Gamma}{eV}\gg\left(\frac{e^2}{\Gamma' C}\right)^2 \tag{B.3}$$

where the coupling is strong enough to extract information about $N(t)$ which is not blurred by feedback and cross-correlation terms (left inequality), but weak enough not to drive the system dot out of equilibrium (right inequality). In this limit the dominating contributions to the detector current's third cumulant are given by $S_3^I \simeq S_3^{I0} + \chi^3 S_3^N$ with

$$\begin{aligned} S_3^{I0} &= \mathcal{T}(1-\mathcal{T})(1-2\mathcal{T})e^4V/h, \\ \chi &= -e^2 d\langle I\rangle/C d\varepsilon', \end{aligned} \tag{B.4}$$

where $\langle I\rangle = \mathcal{T}e^2V/h$ and effective transmission $\mathcal{T} = \Gamma'^2/(\varepsilon'^2 + \Gamma'^2)$. Although the $\sim \chi^3$ term in S_3^I is much smaller than the first one, other terms, corresponding to cross correlations and back action, are negligible compared to the last term.